SOIL TYPE

Citrus may be planted on the coastal clays, coastal peat soils, reef sands, white sands and the interior sands of the intermediate or Rupununi savannahs. These soils have issues of Calcium shortage, Aluminum and iron toxicity, and especially in the case of the clay soils, drainage and compaction issues.

PREPARATION

DRAINAGE

On the clay and pegasse soils drainage is essential. It would be best to plant on cambered beds and drain should be at least 18 inches deep.

On the sands the land is usually sloping and drains along the contour can act as swales to capture rain and irrigation water.

LIMESTONE / ORGANIC MATTER

Limestone is absolutely necessary to:

- Provide calcium
- Reduce acidity
- Promote good soil structure
- Reduce aluminum and iron toxicity due to low pH

Limestone must be ploughed in before planting. On clay soils no more can be ploughed in for the life of the crop. A drench of **AGRICAL** and a regular application of organic matter would help to alleviate the problems of aluminum and iron toxicity and improve soil structure.

For sandy soils limestone applied to the surface can leach down to some extent and can be applied to the surface twice a year. Organic matter would be important for the same reasons as in a clay soil, but in addition would be necessary to hold water and prevent nutrients from leaching.



WEED CONTROL

The approach to weed control is somewhat different for tree crops like citrus. While weeds must be kept under control they protect the soil from erosion. Root exudates from the weeds:

- Improve soil structure
- Release nutrients from the soil
- Increase beneficial soil microbes

If labor is problematic weeds are best controlled with **LIFELINE or CARISTA**. These herbicides would not kill the root and cause minimal damage to the trees due to drift. Slashing has the advantage of maintaining a mulch covering the soil and feeding beneficial soil fauna. Perhaps a good program would include slashing and controlling weeds close to the plants with **LIFELINE or CARISTA**, since slashing near the trees may accidently damage the bark.

SPACING

Spacing should be 12 to 18 feet within the row, and about 25 feet between the rows. This is important to allow a tractor to move between rows for spraying, fertilizing and especially harvesting.

FERTILIZER

The best soil applied fertilizer would be **ELIXIR 7-21-21**. This is a compound fertilizer, which means each grain has all the nutrients. Although it is best to bury the fertilizer **ELIXIR** is still effective when applied to the surface as it dissolves in rain or irrigation water. **FERSAN 15-15-15 or 12-12-17** can be used, but it is blended fertilizer and must be buried.

About 400 lbs. per acre should be applied at planting and at the beginning of each rainy season.

FOLIAR FERTILIZER

Foliar fertilizers are valuable especially at dry times without irrigation, at flowering and to protect roots and stem from Phytopthora.

BUILD A STRONG FOUNDATION.
FOLLOW THE RIGHT PRACTICES.
Better Growth. Better Quality. Better Yields.

Caribbean chemicals
GUYANA LIMITED

"WE SERVE YOUR GROWING NEEDS"

45 Croal Street, Stabroek, G/Town, Guyana
+592-225-4178
exec.guy@caribbeanchem.com
caribbeanchemicals.co.gy

A GUIDE TO CITRUS PRODUCTION



QUALITY PRODUCTS. EXPERT ADVICE.

PEST PROBLEMS

ACOUSHI ANTS

These are best controlled by a bait applied to the trail of ants wherever they are seen. Acoushi ants are especially dangerous to young plants.

HARA HARA

This pest can be controlled by a drench of **PROTECTOR**. Use 15 ml/gal and apply a gallon of the mixture to each tree twice per year.

LEAFMINERS

The citrus leafminer can be controlled by **CAPRID, PROTECTOR, SECURE,** or **CURE.** To control the egg sand young stages use **LEAFGARD.**

PEST PROBLEMS (CONTINUED)

SCALES

These insects can attack the fruit and young stems. They can be controlled by **CAPRID, PROTECTOR, SECURE,** or **CURE.** To control the egg sand young stages use **ADVANCE 10.**

MEALEYBUGS

These insects can attack mostly young stems. They can be controlled by **CAPRID, PROTECTOR, SECURE,** or **CURE.** To control the egg sand young stages use **ADVANCE 10.**

APHIDS

They suck leaves and produce honey dew, which is used by ants for food. They cause a black mold on the leaves because fungi also feed on the honeydew. They can be controlled by **CAPRID, PROTECTOR, SECURE,** or **CURE.** To control the egg sand young stages use **ADVANCE 10.**

MITES

They can cause damage to leaves and fruit. They can be controlled by **SECURE,** or **CURE.** To control the egg sand young stages use **AZTEZ.**

DISEASE PROBLEMS

GUMMOSIS / PHYTOPTHORA	Best controlled by **PHORTIFY, QUAD 14, PILOT** or **ACROBAT.**
SCAB AND OTHER DISEASES AFFECTING APPEARANCE	Best controlled by **CUPROFIX, GLORY.**
NEMATODES	Attack roots in sandy soil.





Caribbean chemicals GUYANA LIMITED

"WE SERVE YOUR GROWING NEEDS"

PATHOGEN CONTROL PROGRAM

Twice per year drench each tree with **PROTECTOR** at 15gm/gal. Use a gallon of mix per tree.
On sands add **VYD-8** at 10 ml/gal to the mix.

AS ROUTINE THROUGH THE LIFE OF THE CROP:

ALTERNATE THE FOLLOWING 2 SPRAY MIXTURES EVERY 2 WEEKS

MIXTURE 1		MIXTURE 2	
SECURE	5ml/gal	PROTECTOR	5gm/gal
LEAFGARD	5gm/gal	ADVANCE 10	10ml/gal
PHORTIFY	10ml/gal	K41	10ml/gal
CALMAX	10ml/gal	BIO 20	10ml/gal
PILOT	10gm/gal	CUPROFIX	10ml/gal
NUFILM	2ml/gal	NUFILM	2ml/gal

Consistent monitoring and correct application of these products will help maintain healthy trees and high yields.



QUALITY PRODUCTS. EXPERT ADVICE.
Your Partner in Successful Farming.