Group 5 - Root & Tuber Crops

These crops are grown from cuttings, not seeds, so they can handle soil-applied Moccasin 96 EC — as long as they are planted deep enough.



Sweet Potatoes

- Apply before planting
- Plant deep — at least 3 nodes below the soil surface
- After planting: apply between rows after 7 days



Cassava

- Apply 2 to 3 days before planting
- Can spray over the top immediately after planting only
- Do NOT spray once young leaves begin to emerge

NUTGRASS? Moccasin 96 EC kills nutgrass leaves as they emerge. For best results, plough first to damage the tubers, then apply.

BEFORE YOU SPRAY KNOW THESE RULES

No Surfactant

Never add a surfactant, crop oil, or wetting agent to the tank. It will force the chemical through the leaf and damage your crop.

Needs Moisture

Apply when the soil is damp, or irrigate lightly after spraying. Dry soil means the chemical cannot form its weed barrier.

Sandy Soils

Reduce your rate on coarse or sandy soils. Moccasin 96 EC leaches downward quickly and can reach crop roots.

Rate of Application
20ml (4 tablespoons) per gallon



We Serve Your Growing Needs
Caribbean Chemicals Guyana Ltd
45 Croal Street, Stabroek, Georgetown, Guyana

+592 225-4178
+592 225-9639

Message on Whats App
+592 743-4769

UPL | Caribbean chemicals GUYANA LIMITED



MOCCASIN 96 EC

Reliable Residual Weed Control
For Stronger Crop Establishment

UPL | Caribbean chemicals GUYANA LIMITED

Moccasin 96 EC works in the top layer of soil. It targets weeds as they germinate — before they emerge. Your job is to apply it correctly so it catches the weed, not your crop.

GROUP 1 - Safe to Spray Over the Top

These crops are strong enough to handle direct application once they are established.



Corn

- Plant your seed and apply
- Make sure seed is properly buried before spraying
- Can be used while crop is still growing — up to 40 inches tall



Cabbage / Pakchoy — Tomato / Pepper

- Before transplanting: Apply 1 to 3 days before you put plants in the ground
- After transplanting: Wait 7 to 10 days until roots and leaves are firm and hardened
- Do not spray over the top if plants are stressed, wet, or under hot sun.

Group 2 - Soil Only. Do Not Spray the Plant

These crops can handle the chemical in the soil, but their leaves are too delicate for direct spraying.



Eggplant

- Do NOT spray over the top — ever
- Apply only between the rows using a shielded sprayer



Celery & Lettuce

- Do NOT spray over the top
- Apply before transplanting only
- Make sure roots are set well below the soil surface
- When transplanting, do not let surface soil

Group 3 - Not Recommended

Okra, Bora, Watermelon, Cucumber, Squash, Caraille. These crops are too sensitive. Risk of stem damage and crop loss is high. Avoid use or consult your CCG representative before applying.

Group 4 - Perennial Crops & Fruit Trees

Apply to the soil. Keep spray away from the plant at all times.



Passion Fruit

- Apply to bare soil only
- Do NOT let spray touch the leaves



Banana, Plantain & Papaya

- Apply to bare soil only
- Do not let spray touch young stems
- For papaya: Wait 14 days after transplant-



Mature Fruit Trees

- Apply beneath the tree
- Keep spray away from fruit and leaves